

Published in final edited form as:

Clin Child Fam Psychol Rev. 2014 December ; 17(4): 399–411. doi:10.1007/s10567-014-0174-2.

Posttraumatic stress following acute medical trauma in children: A proposed model of bio-psycho-social processes during the peri-trauma period

Meghan L. Marsac, PhD,

The Center for Injury Research and Prevention, The Children's Hospital of Philadelphia,
Department of Psychiatry, University of Pennsylvania

Nancy Kassam-Adams, PhD,

The Center for Injury Research and Prevention, The Children's Hospital of Philadelphia;
Department of Pediatrics, University of Pennsylvania

Douglas L. Delahanty, PhD,

Department of Psychology, Kent State University

Keith Widaman, PhD, and

Department of Psychology, University of California-Davis

Lamia P. Barakat, PhD

Division of Oncology, The Children's Hospital of Philadelphia

Department of Pediatrics, University of Pennsylvania

Abstract

Millions of children worldwide experience acute medical events. Children's responses to these events range from transient distress to significant posttraumatic stress disorder symptoms (PTSS). While many models suggest explanations for the development and maintenance of PTSS in adults, very few have focused on children. Current models of child PTSS are primarily restricted to the post-trauma period, thus neglecting the critical peri-trauma period when screening and preventive interventions may be most easily implemented. Research on PTSS in response to pediatric medical trauma typically examines predictors in isolation, often overlooking potentially important interactions. This paper proposes a new model utilizing the bio-psycho-social framework and focusing on peri-trauma processes of acute medical events. Understanding the relationships among bio-psycho-social factors during peri-trauma can inform early identification of at-risk children, preventive interventions, and clinical care. Recommendations for future research, including the need to examine PTSS in the context of multiple influences, are discussed.

Corresponding author: Meghan L. Marsac, Center for Injury Research and Prevention, The Children's Hospital of Philadelphia, 3535 Market, Suite 1150, Philadelphia, PA 19104, USA. Telephone: 267-426-9620. marsac@email.chop.edu.

Conflicts of Interest

The authors declare that they have no conflict of interest.

Keywords

child PTSD; child PTSS; medical traumatic stress; peri-trauma; acute medical trauma; bio-psycho-social model

Events related to injury, acute medical illness, and medical treatment are among the most common potentially traumatic events in childhood (Murray & Lopez, 1996). In the United States alone, twenty million children suffer unintentional injuries each year (Grossman, 2000), with many more children facing sudden illnesses (Marks & McQueen, 2001). An acute medical event is defined as a sudden, unexpected, and new medical event for a child. This includes injuries or sudden illnesses (excluding chronic illnesses). Unique to medical trauma, the peri-trauma period typically includes physical trauma (i.e., discomfort and/or pain due to injury, illness, or medical treatment), interaction with medical treatment providers, and medical interventions; all components of this experience can contribute to the child's emotional reactions to his or her injury or illness. The peri-trauma period in children exposed to acute medical events is also unique from other acute trauma in presenting the opportunity for implementation of systematic screening and preventive interventions while children are interfacing with the medical system for physical care. Children facing acute pediatric medical events (e.g., injury, ICU admission) frequently exhibit significant posttraumatic stress symptoms (Balluffi et al., 2004; Kean, Kelsay, Wamboldt, & Wamboldt, 2006; Mintzer et al., 2005; Walker, Harris, Baker, Kelly, & Houghton, 1999). More specifically, a meta-analysis found that 19% of children with injuries and 12% of children with illnesses develop persistent PTSS (Kahana, Feeny, Youngstrom, & Drotar, 2006). The consequences of PTSS extend beyond the psychological symptoms to also include poorer functional health outcomes and greater use of health services (Holbrook et al., 2005; Landolt, Vollrath, Gnehm, & Sennhauser, 2009; Mintzer, et al., 2005; Zatzick et al., 2008). Thus, the major impact of PTSS on children demands clinically-relevant research to support recovery after potentially traumatic acute medical events.

In general, posttraumatic stress disorder (PTSD) requires exposure to a potentially traumatic event (i.e., experiencing an event which involved the possibility of death or serious injury to the individual or to others) and includes symptoms of re-experiencing or intrusion, avoidance, arousal, changes in mood or cognitions, and impaired daily functioning (American Psychiatric Association, 1994; 2010). For both clinical and research purposes, it is important to note that significant PTSS, regardless of PTSD diagnostic status, are associated with negative outcomes (Holbrook, et al., 2005; Stoddard & Saxe, 2001). In children with PTSS related to medical events, re-experiencing can take the form of dreams (e.g., nightmares about a car crash) or can be displayed in play (e.g., drawing pictures of a surgery). Arousal can be demonstrated as an increased startle response or activation of physiological responses following a trauma trigger (e.g., heart rate elevated at a follow-up appointment). Avoidance can be observed in a variety of ways such as refusing to take medications, not wanting to talk about the injury or illness, or refusing to return to the hospital. With the new DSM-5 criteria, prevalence rates for PTSD in adult trauma victims are expected to remain the same (Calhoun et al., 2012). It is unclear how the prevalence of PTSD/PTSS will change in children. However, given that depressive symptoms can also be

a result of exposure to acute trauma in children (Kassam-Adams, Marsac, & Cirilli, 2010), it is possible that the changes to DSM-5 regarding mood criteria will result in higher rates of PTSD/PTSS in children exposed to acute medical events.

Specific to PTSS due to medical events, Kazak and colleagues (2006) explained that families are best served when medical teams apply an integrative model to understanding, preventing, and treating medical traumatic stress. Their proposed framework highlights the importance of recognizing the systems (e.g., school) with which the child and family interact and examining children's responses to the medical event using a trauma phase approach (pre-, peri-, post-). Research should include information on all phases of the medical trauma and consider developmental context and environmental influences (Kazak et al., 2006).

In children, research has focused on the post-trauma phase of acute events. Data during the peri-trauma period is particularly sparse. In acute medical trauma, the peri-trauma period can be defined as the time from which the potentially traumatic event (e.g., a child is injured in a motor vehicle crash) begins through the child's medical treatment (i.e., including the medical treatment as part of the potentially traumatic event such as emergency transport, surgery, and/or hospitalization). In medical populations, peri-trauma factors are of special importance. Applying a bio-psycho-social framework to the prevention of PTSS, biological markers during peri-trauma have the potential to serve as readily available screeners, while psycho-social factors can serve as potential mechanisms of action to prevent or reduce sustained PTSS. Understanding markers and mechanism of action provides a unique opportunity for prevention of PTSS during the child's medical treatment.

To date, preventive PTSS interventions for pediatric medical populations have produced mixed results (Berkowitz, Stover, & Marans, 2011; Cox & Kenardy, 2010; Kramer & Landolt, 2011; Stallard et al., 2006; Zehnder, Meuli, & Landolt, 2010). For example, two preventive interventions have shown promise in preventing PTSS during the post-trauma period following acute medical events: Kids and Accidents (psycho-education with a web-based component for children and a booklet for parents) and the Child and Family Traumatic Stress Intervention (four-session in-person psycho-educational and skills based intervention) (Cox & Kenardy 2010; Berkowitz, Stover, & Marans, 2011). In contrast, though not specific to pediatric medical events, some efforts to prevent during the peri-trauma or early post-trauma period (e.g. via debriefing) have produced no effect or worsened symptoms by interfering with the normal recovery process (McNally, Bryant, & Ehlers, 2003; Rose, Bisson, Churchill, & Wessely, 2002). Given these results, it is unclear what intervention components or mechanisms of action are most effective and what factors should be targeted in developing or improving preventive interventions. In addition, most children exposed to acute medical events do not have contact with behavioral health providers; thus, screening and intervening by medical providers during the peri-trauma period has the potential to significantly extend the reach of secondary prevention programs. Recognizing and working to prevent PTSS in the peri-trauma period can help improve children's care, promoting full emotional and physical recovery.

This paper provides a narrative review, synthesizing current research and theory regarding the development and maintenance of PTSS in children exposed to acute medical events.

Extending current theory, we propose a new model that integrates key biological, psychological, and social processes during the peri-trauma phase that may contribute to the development and maintenance of PTSS. Specifically, we summarize factors across each trauma phase (pre-, peri-, post-) using the biological, psychological, and social/environmental framework. We focus the discussion on the peri-trauma time period, given its importance in acute medical trauma and current gaps in the literature. See Fig. 1 for a summary of the relationships in the proposed model.

Current Theoretical Models of PTSS/PTSD Development in Children

Below we provide a brief history of the bio-psycho-social framework and three primary theories which influenced the model proposed in this paper.

Bio-psycho-social framework

As understanding of psychological disorders and symptoms has progressed, it has become increasingly clear that the onset of symptoms does not have a single cause. Rather, psychological symptoms should be assessed and treated within the context of the individual and his or her situation. Engel proposed using a bio-psycho-social framework to guide medicine (including psychiatry) in the 1970s (Engel, 1977). Thirty years later, Fava and Sonino (2008) summarized how the bio-psycho-social framework has been applied to patient care. They noted that while the evidence base for the bio-psycho-social framework has grown over time, actual implementation of this framework in patient care has been slow (Fava & Sonino, 2008). In the field of psychology, Suls and Rothman (2004) suggested that the application of bio-psycho-social framework has yet to be utilized to its full potential and should be considered in emerging theories of health psychology and clinical practice. Research has identified biological, psychological, and social (environmental) contributors to the onset and persistence of PTSD/PTSS (Alisic, Jongmans, van Wesel, & Kleber, 2011; American Psychiatric Association, 2014; Cox, Kenardy, & Hendrikz, 2008; Kahana, et al., 2006), and using the bio-psycho-social framework to inform treatment of PTSD/PTSS has been suggested (Finkelhor, Turner, Ormrod, & Hamby, 2009). Despite current research and recommendations for clinical practice, the bio-psycho-social framework has yet to be described clearly as it relates to the conceptualization of the development of PTSD/PTSS and has not been integrated into standard clinical care.

Biological Model

Schwartz and Perry (1994) suggested that neurobiology plays a primary role in the development of PTSS: Physiological symptoms maintain hyper-arousal and influence the formation of memories. When faced with an acute potentially traumatic stressful event, the body's natural reaction is to activate systems which create the fight-or-flight response. As this response is initiated, changes in the brain occur to promote survival. In some individuals, this response continues beyond the duration of the stressor or at a higher frequency or intensity. Thus, rather than the fight-or-flight response enhancing survival as intended, the response may create PTSS (Schwartz & Perry, 1994). Schwartz and Perry suggest that several factors contribute to this process: genetics, family factors, the developing brain (i.e., how developed the brain is when a trauma exposure occurs), and

malignant memories. Genetics play a role in how a child develops arousal symptoms and how a child learns; thus, genetic predisposition indirectly impacts the development of PTSD. The family environment can impact how the child processes traumatic events, and memories have a stronger impact on the developing brain than the adult brain. During periods of high stress, neurobiological processes can disrupt the process of memory consolidation resulting in the formation of “malignant memories”, which are related to the later development of trauma symptoms (particularly those related to arousal; Schwartz & Perry, 1994). Though not specific to acute pediatric medical events, research supports the role of malignant memories in that heightened peri-trauma physiological responses (e.g., elevated heart rate, release of hormones) are associated with aberrant memory function (Pitman & Delahanty, 2005; Pitman et al., 2012).

Schwartz and Perry (1994) were among the first to highlight how physiological changes in the brain interact with memory/cognition and the child’s social environment to affect the development and maintenance of PTSD. While they allude to the interactions among these specific biological, psychological, and environmental processes, they place heavy emphasis on the biological processes involved in memory formation and neglect more specific child and parent psychological processes (e.g., appraisals, coping behavior). This theory focuses on the post-trauma phase and chronic PTSD, rather than processes occurring during the peri-trauma phase, restricting the extent to which this theory can inform preventive efforts.

Psychological Model

A model proposed by Meiser-Stedman (2002) suggests the application of cognitive behavioral theory (CBT) toward understanding and treating PTSD in children. His rationale highlights the role of children’s cognition and memory in the maintenance of PTSS. Meiser-Stedman offers support for a CBT model based on modifications of theories offered to explain the maintenance of PTSD in adults. He suggests that this allows for 1) consideration of a developmental framework, 2) inclusion of the role of the family, 3) examination of the child’s pre-trauma functioning, 4) attention to the increase in comorbid psychological disorders, 5) examination of the child’s ability to verbalize memories and cognitions, and 6) consideration of the role of maladaptive cognitions, particularly cognitive avoidance. Meiser-Stedman and colleagues (2007; 2009) present evidence to support the theory that children’s memories and cognitions play a role in the maintenance of PTSS (Meiser-Stedman, Dalgleish, Glucksman, Yule, & Smith, 2009; Meiser-Stedman, Smith, Yule, Dalgleish, & Glucksman, 2007). More specifically, in children exposed to physical assaults or motor vehicle accidents, maladaptive appraisals mediated the relationship between initial PTSS and PTSS six months later (Meiser-Stedman, et al., 2009).

Meiser-Stedman’s CBT model has positively influenced the field by encouraging researchers and clinicians to recognize similarities and differences in PTSS maintenance among adults and children and by bringing attention to the importance of children’s cognitions. This model suggests consideration for psychological factors (e.g., pre-, post-trauma symptoms) and the environment (i.e., family roles). Yet, the model does not account for variability in biological factors nor does it include child and parent coping behaviors.

Further, his model focuses on the pre- and post-trauma phases, again leaving out the potentially critical peri-trauma phase.

Social/environmental model

Finally, Kazak's (2006) application of the Pediatric Psychosocial Preventative Health Model (PPPHM) for assessment and treatment of children in health care settings reinforces the concept of a bio-psycho-social model for preventive health. She brings together the influence of social factors on children by using the PPPHM. The PPPHM highlights the bi-directional influence between a child's health and their multi-layered systems (e.g., medical, school, and family systems). For example, a child's illness may affect parenting techniques and school attendance, which then, in turn, affects the child. Kazak suggests that this model can be particularly helpful in determining the level of preventive care needed to facilitate a child's recovery. Research has supported using the PPPHM to determine child risk and level of care needed (Kazak, 2006).

While PPPHM reinforces the idea of implementing the bio-psycho-social framework for PTSS related to medical events and highlights the importance of a child's social environment, it does not clearly delineate specific factors that may be important during each trauma phase. Because this model focuses more generally on preventive health rather than specifically on PTSD/PTSS, specific factors associated with PTSD/PTSS are not discussed (e.g., trauma history, appraisals).

Summary

Taken together, these three models support the concept of applying the bio-psycho-social framework to the development of PTSD/PTSS in children exposed to acute medical events. While drawing from each of these models, the current proposed model is innovative in several ways. The proposed model 1) brings together bio-psycho-social factors and their interaction into a single overarching model, 2) focuses on pediatric acute medical trauma, and 3) highlights the importance of the peri-trauma phase in medical events.

Proposed Model and Supporting Evidence

Overview

Influenced by the models above and based in evidence described below, we propose a model which applies a bio-psycho-social framework to the development of PTSD/PTSS in children experiencing acute medical trauma. The proposed model suggests factors across each trauma phase, with particular focus on the peri-trauma phase. As noted, the peri-trauma phase of acute medical trauma may be an ideal time for screening and intervention and has often been neglected in current research and practice. See Figure 1 for an illustration of the proposed relationships between variables in the model.

During the pre-trauma phase of the event, specific biological factors (child genetic predisposition, sex, age), psychological factors (emotional functioning), and social factors (child trauma history, family functioning, community support, culture) ought to be considered in determining risk for subsequent PTSS to allocate resources to those likely to

have the most need. In addition to contributing directly to PTSS, pre-trauma variables also contribute to the processes during the peri-trauma period (Fig.1, path a). For example, girls report more early PTSS, and parents of younger children use more coping assistance (Gold, Kant, & Kim, 2008; Marsac, Donlon, Winston, & Kassam-Adams, 2013). While we do not provide an exhaustive list of specific potential relationships between pre- and peri-trauma variables, we suggest these relationships should be considered in future research. During the peri-trauma period, the intra- and inter-relationships among biological (initial memory formation, hormone response, cardiovascular reactions), psychological (early child PTSS, appraisals, coping), and social/environmental (trauma severity, medical team support, community support, parent PTSS, parent appraisals, parent coping assistance) may affect the development and maintenance of PTSD/PTSS. To date, most research has focused these variables in isolation of one another, particularly during the peri-trauma period. Table 2 provides a summary of the evidence to date to support the independent role of each of these variables in contributing to PTSS. In addition to the independent contributions of these factors, we suggest examining these variables together as described below (see Peri-Trauma Period: Integrating Biological, Psychological, and Social Factors). During post-trauma, factors contributing to PTSS include biological (physical injury recovery), psychological (non-PTSS emotional reactions), and social (parent coping assistance, community support) components. Again, in addition to their independent contribution to PTSS, pre and peri-trauma variables may influence these factors (Fig. 1, path o). For example, physical recovery from injury is associated with pre-existing emotional health, child gender (females at risk for worse recovery) and severity of physical trauma (Polinder et al., 2005; Zonfrillo, Durbin, Winston, Zhao, & Stineman, 2013). While it is beyond the scope of this paper to detail every potential relationship across time points, we include these post-trauma factors to encourage future research to improve understanding of the development of PTSS and to increase screening and intervention effectiveness.

Peri-Trauma Period: Biological Factors and PTSS

Acute physiological responses, including elevated stress hormone levels and heart rate, may serve as early predictors of PTSS in children. Cortisol levels have been examined in PTSD research, although the direction of alterations in cortisol related to PTSD/PTSS outcomes have been mixed in adults (Klaassens, Giltay, Cuijpers, van Veen, & Zitman, 2012; Meewisse, Reitsma, de Vries, Gersons, & Olff, 2007). Individual differences in the symptom presentation of PTSD and time since trauma may serve to explain these variations in findings between cortisol levels and PTSD (Miller, Wolfe, Fabricant, & Stein, 2009; Weems & Carrion, 2007); however, the effect of time has only been demonstrated during the post-trauma period (i.e., comparing children with trauma within the past year to those with trauma more than a year ago; Weems & Carrion, 2007). In children, there has been greater consistency regarding the direction of the relationship between early peri-trauma stress hormone levels and PTSD, especially in boys (Carrion et al., 2002; Delahanty, Nugent, Christopher, & Walsh, 2005; Ostrowski, Christopher, van Dulmen, & Delahanty, 2007). For example, Ostrowski and colleagues (2007) assessed urinary cortisol levels in 54 children admitted for injuries at baseline, six-weeks, and seven-months. At baseline, levels of stress hormones were measured in urine samples with collection initiating upon

admittance to the hospital. PTSS was assessed at six weeks and seven months. At both follow-ups, higher cortisol levels predicted PTSS only in boys.

Perhaps the most consistently observed biological risk factor for PTSD is heart rate assessed within the first 24 hours of hospital admission for injury. In-hospital heart rate levels have been found to predict PTSD at six weeks (Nugent, Christopher, & Delahanty, 2006b), three months (Kassam-Adams, Garcia-Espana, Fein, & Winston, 2005), and six months post-injury (Bryant, Salmon, Sinclair, & Davidson, 2007a; De Young, Kenardy, & Spence, 2007; Nugent, et al., 2006b), even after controlling for age, sex, and injury severity. A single measurement (earliest available heart rate) has been shown to be a stronger predictor of PTSD than heart rate averaged over the first 20 minutes post-admission (Nugent, Christopher, & Delahanty, 2006a). For example, Nugent and colleagues (2006a) collected heart rate for 82 children who were admitted to the emergency department for treatment of an injury, and collected PTSS data at six weeks and six months. Heart rate was collected during EMS transport, upon arrival to the emergency department, for the first 20 minutes of the emergency department visit, and at discharge. After controlling for demographic factors and depressive symptoms, the first recorded heart rate and heart rate averaged over the first 20 minutes following admission both predicted PTSS at six weeks; however, only the first recorded heart rate predicted PTSS at six months (Nugent, et al., 2006b). The role of heart rate in medical trauma other than injury is more complicated due to the potential effects of medication and disease processes and warrants further investigation. Heart rate upon admission has potential public health relevance as a readily available and easily implemented screener of PTSD/PTSS risk, requiring no additional procedures. Mechanisms through which heart rate is associated with PTSD/PTSS remain unclear, though it is suggested that when an individual appraises the situation as more stressful, he or she has stronger physiological responses (e.g., increased heart rate; Fig. 1, path d; Olff, Langeland, & Gersons, 2005).

Peri-Trauma Period: Psychological Factors and PTSS

Folkman and Lazarus (1985; 1991) were the first to suggest an essential role for cognitive appraisals in evaluating an individual's coping responses to stress (Folkman & Lazarus, 1985; Lazarus, 1991), positing that a primary appraisal occurs when the individual interprets the event as threatening. Following primary appraisal, the individual selects a coping technique (Folkman & Lazarus, 1985). Drawing from literature examining the manifestation of anxiety, maladaptive cognitive appraisals (e.g., nothing will be normal again) after a difficult event lead to behavioral strategies (i.e., coping) that directly produce PTSS and/or prevent the development of realistic and adaptive appraisals (Ehlers & Clark, 2000; Meiser-Stedman, 2002). Though these theories were not posed specifically to pediatric medical trauma, they can be applied. For example, after a frightening medical event, a child may be naturally exposed to potential trauma triggers (e.g., follow-up appointments requiring return to the hospital). This exposure to a reminder of the event can promote realistic appraisals and help to diminish emotional distress. However, if a child has appraised their injury or illness as particularly traumatic, this may lead to avoidant coping strategies (Fig. 1, path f). Avoidant coping may then reduce exposure to reminders and divert the natural recovery process.

Research has supported the independent roles of appraisals and coping in children facing acute medical trauma. Specific to appraisals, children's cognitions (e.g., perception of threat during the traumatic event, negative appraisals about vulnerability to future harm, and negative interpretation of intrusive memories, alienation from others, anger, and rumination) are related to worse PTSS (Bryant, Salmon, Sinclair, & Davidson, 2007b; Ehlers, Mayou, & Bryant, 2003; McDermott & Cvitanovich, 2000; Stallard & Smith, 2007). Amount and type of child coping with acute medical events post-trauma are also related to PTSS, though peri-trauma coping and PTSS have yet to be examined (Alisic, Conroy, Magyar, Babl, & O'Donnell; Marsac, Donlon, Hildenbrand, Winston, & Kassam-Adams, 2014; Stallard & Smith, 2007; Zehnder, Prchal, Vollrath, & Landolt, 2006). In a study of 243 children with injuries, Marsac and colleagues (2014) found that worse PTSS during the peri-trauma period was associated with later use of distraction, social withdrawal, problem-solving, and blaming others (Marsac, et al., 2014). In children with newly diagnosed diabetes ($n = 38$) or an injury or newly diagnosed chronic disease ($n = 161$), use of active coping and support-seeking coping was weakly related to PTSS, and use of religious coping predicted fewer PTSS one year later (Landolt, Vollrath, & Ribi, 2002; Zehnder, et al., 2006). These studies suggest a probable role for the interaction among peri-trauma PTSS, coping, and appraisals in predicting later PTSS. However, more research is necessary to better understand these relationships.

Peri-Trauma Period: Social (Environmental) Factors and PTSS

Findings for the relationship between severity of trauma exposure for acute medical events and later PTSS has been mixed, ranging from no effect to large effect sizes (Alisic, et al., 2011; Cox, et al., 2008; Kahana, et al., 2006; Trickey, Siddaway, Meiser-Stedman, Serpell, & Field, 2012). More research is needed to understand if the differences in findings could be explained by other peri-trauma factors serving as moderators. While support from medical teams (e.g., delivering trauma-informed care) and support from the community has been recommended, empirical evidence is not available on the impact of this type of support during the peri-trauma period (Kassam-Adams, Marsac, Hildenbrand, & Winston, 2013; Kazak, et al., 2006). Based on research during the post-trauma period, we expect community support (including the medical community) during the peri-trauma period could also help prevent PTSD/PTSS (Kahana, et al., 2006; Trickey, et al., 2012).

Parents may influence the development of PTSD/PTSS through their own PTSS, cognitive appraisals, and efforts to help their child cope (i.e., coping assistance; Barrett, Dadds, & Rapee, 1996; R Blount, Davis, Powers, & Roberts, 1991; Power, 2004; Salmon & Bryant, 2002; Skinner & Zimmer-Gembeck, 2007). A number of studies have identified a relationship between caregiver and child PTSS during the post-trauma period (Alisic, et al., 2011; Cox, et al., 2008; Trickey, et al., 2012). While there has not yet been a focused effort to examine parents' distress during the peri-trauma period, Balluffi et al. (2004) found a relationship between parents' PTSS during their child's intensive care hospitalization and their child's PTSS two months later. Additionally, a recent study conducted by De Young and colleagues (2014) identified an association between child and parent PTSS one and six months following a burn injury. Though neither study (i.e., Balluffi et al., De Young et al.) assessed for a potential association between peri-trauma child PTSS and peri-trauma parent

PTSS, combining these results with others in the post-trauma phase suggests a potential relationship between peri-trauma parent PTSS and subsequent child PTSS (Alisic, et al., 2011; Balluffi, et al., 2004; Cox, et al., 2008; Trickey, et al., 2012).

Several studies have found that in children with PTSS related to medical events, early child PTSS predicts later PTSS (Dagleish et al., 2008; Kassam-Adams & Winston, 2004). For example, Kassam-Adams and Winston (2004) conducted a study to examine PTSS within 1 month following pediatric injury and again 3 months later (sample $n = 243$). Findings indicated that while a diagnosis of acute stress disorder was not the best predictor of posttraumatic stress disorder, a significant relationship emerged between PTSS severity ($r = .56, p < .0005$) (Kassam-Adams & Winston, 2004). Similarly, Dagleish and colleagues (2008) followed 367 children (ages 6-17 years old) for six months following an injury related to a motor vehicle crash; results indicated that PTSS two to four weeks post-injury significantly predicted PTSD six months later (Dagleish, et al., 2008). No research to date suggests that parent appraisals or coping assistance has a direct relationship on child PTSS. These factors are potential mechanisms through which parents may influence their child's PTSS and are described below.

Peri-Trauma Period: Integrating Biological, Psychological, and Social Factors

Very little empirical evidence exists to help explain the interactions among the bio-psycho-social factors addressed in this manuscript and how, when integrated, they may influence the onset or severity of child PTSS. Thus, the model proposed here aims to fuse together theory and empirical evidence above and highlight areas for new research to inform screening and preventive intervention.

As described above, physiological arousal is related to PTSS outcomes. Integrating this knowledge with the model of how biological, psychological, and social processes contribute to PTSS, we can speculate on the nature of these relationships and examine the intra-relationships of the peri-trauma variables. For example, hormonal responses and cardiovascular levels are related to the formation of memories (Fig. 1, path b), which may in turn influence PTSS. Heart rate is also related to the psychological variable of appraisals (path d) (Nixon et al., 2010; Pitman & Delahanty, 2005; Pitman, et al., 2012); both heart rate and appraisals are related to PTSS (Meiser-Stedman, et al., 2009; Nugent, et al., 2006a). Thus, there may be a shared variance or moderated affect, with appraisals influencing heart rate (path d). Changing appraisals during peri-trauma (e.g., using cognitive behavioral techniques) may serve to also affect heart rate and memory formation. Elevated heart rate could serve as a screener to identify intervention need.

However, appraisals do not operate in isolation: appraisals are related both to psychological variables of early PTSS (Fig. 1, path e) and coping (path f). Specifically, how children perceive an event while it is happening predicts PTSS (Ehlers, Mayou, & Bryant, 2003; McDermott & Cvitanovich, 2000). Extrapolating from research during the post-trauma period, in a study conducted by Stallard and colleagues (2007) of cognitive appraisals and coping with concurrent PTSS in injured children ($N = 75$), appraisals and coping together accounted for 64% of the variance in PTSS 8 months after injury. Interestingly, trauma memory did not significantly contributed to PTSS in this study, suggesting that the

relationships between appraisals and memory (path c) need further exploration (Stallard & Smith, 2007). Interventions during the peri-trauma period which focus on appraisals and coping together could be more powerful than those aiming to change appraisals only. It is also possible that the relationship between appraisals and PTSS is partially accounted for by coping behaviors.

We propose that appraisals during the peri-trauma period are also influenced by the social variable parent appraisals (Fig. 1, path k) and coping by parent coping assistance (path m). Parental influence over child appraisals and coping has been most clearly established as related to child anxiety. Drawing from the larger appraisals and coping literature (i.e., beyond pediatric medical events and PTSS), direct observation of parent-child processes highlight that parents' role in how children appraise situations. Parents of children with anxiety indicate that they have more threatening appraisals of events and maladaptive coping strategies than parents of children without anxiety (Barrett, Rapee, Dadds, & Ryan, 1996; Micco & Ehrenreich, 2008). For example, Barrett and colleagues (1996) presented a series of hypothetical scenarios separately to 205 children and parents to elicit cognitive appraisals (threat versus neutral interpretations of ambiguous situations) and coping responses (adaptive versus maladaptive). In a subsequent family discussion task using these situations, anxious children and their parents generated more threat appraisals and more avoidant coping responses than non-anxious participants (Barrett, Rapee, et al., 1996). Further, parents of anxious children were more likely to reciprocate and reward avoidant coping suggestions made by their children. When parents encouraged avoidant coping, children were more likely to sustain these strategies (Dadds, Barrett, Rapee, & Ryan, 1996). Specific to children exposed to trauma, parent coping assistance has been associated with child coping for community violence (Kliewer et al., 2006), pediatric injury (Marsac, et al., 2014), and painful medical procedures (R Blount et al., 1989). For example, injured children whose parents engaged in more coping assistance strategies used more social support as a coping strategy (Marsac, et al., 2014). Evidence suggests a potential role for parent influence over appraisals and coping during the peri-trauma period (paths k, m), but investigations are warranted to clarify this role. Further understanding how children's appraisals develop by taking each factor into consideration has direct implications for preventive interventions and clinical care. For example, if parent appraisals strongly influence child appraisals, then parent appraisals can be targeted with cognitive behavioral treatment as well.

Psychological variables of early child PTSS and child coping are also influenced by the child's social environment (Fig. 1, paths g, i) such as trauma severity, support from the medical team, community support, and parent PTSS (Alisic, et al., 2011; Balluffi, et al., 2004; Cox, et al., 2008; Kahana, et al., 2006; Kazak, et al., 2006; Trickey, et al., 2012). As mentioned previously, the relationship on trauma severity to PTSS has ranged from no effect to a large effect across samples (Alisic, et al., 2011; Cox, et al., 2008; Kahana, et al., 2006; Trickey, et al., 2012). However, the reason in this variation is not well understood. It could be that some of the other peri-trauma variables are impacting this relationship. Additionally, to date, while it is recommended that medical teams deliver trauma-informed care, the effect of doing so is unknown (path g). Community support (e.g., peer, school) has been shown as a protective factor in the post-trauma period but has yet to be examined in the peri-trauma period (Kahana, et al., 2006; Trickey, et al., 2012), so the intra- peri-trauma period

relationships are unclear (path g). Finally, the relationship between parent and child PTSS is also complex (path i). For example, Ostrowski et al. evaluated 118 children admitted to hospital for an injury. Child avoidance symptoms in the hospital predicted PTSS two weeks later; an interaction between child and parent PTSS at two weeks predicted child six-week PTSS. Thus, the interaction between child and parent PTSS may be important in examining longer term outcomes (Ostrowski et al., 2011). It may be that if the medical team supports the parent and addresses parental symptoms, the parent may be better able to support the child's coping.

Limitations

A goal of proposing this new model is to stimulate future research focusing in on the peri-trauma time period of pediatric medical events that applies a bio-psycho-social framework. Thus, the current evidence to date to support this model is limited and recommendations for clinical practice should be interpreted with caution. In addition, while biological markers of heart rate and cortisol levels have the potential to be non-invasive and easily obtainable, currently there is not a simple way for a clinician to "score" them, given individual variability. In the future, new technology may make this possible. In the evidence reviewed above, many of the studies identified findings with small effect sizes, which bring into question the confidence that we can have that these findings have significant clinical relevance. Additionally, with small effect sizes, it is difficult to determine the utility of individual risk factors for screening and intervention. Further, no study has examined the all (or most) of the relationships proposed in the model together. When these factors are brought together, the intra- and inter-relationships may change. This review also limited its scope to acute medical trauma rather than including all acute trauma; future work should examine how the model proposed here could be applied across acute trauma. Specifically, future research can highlight the similarities and differences in the bio-psycho-social variables across trauma types. Finally, to date there is a paucity of research on the cross-cultural applications of these specific variables in relation to PTSD/PTSS. More research is needed to support cross-cultural applications and/or need adaptations.

Summary

The need to support children exposed to acute medical events is clear, with millions of children worldwide experiencing acute medical events every year (Murray & Lopez, 1996). Over the past thirty years, the field has significantly advanced our understanding of how the presentation of PTSD/PTSS in children differs from adults. However, ideal screening to predict sustained PTSS and efforts at preventive interventions are in their infancy and offer mixed results. The development and refinement of these preventive interventions can benefit from a better understanding of processes during the peri-trauma phase. Current models of PTSD/PTSS provide some guidance for factors to target moving forward; however, none of the current models of the development of child PTSD/PTSS bring together biological, psychological, and social forces and their interactions. Further, to date, no models have focused on the peri-trauma period when screening and interventions can be delivered as part of medical care. The current model serves to fill both of these gaps and stimulate additional

research to advance our understanding of the development of PTSS in children following acute medical trauma.

Adopting a bio-psycho-social model of PTSD/PTSS development and maintenance in children can serve to guide future research. Specifically, future research will significantly move the field forward by focusing on two main areas: 1) the intra- and inter-relationships among bio-psycho-social factors associated with PTSD/PTSS and how these relationships predict long-term outcomes in children, and 2) the processes involved during the peri-trauma phase. An important area of future research related to the inter-relationships among bio-psycho-social factors includes exploration of how peri-trauma physiological responses (bio) are related to early appraisals (psycho) and whether parental behavior (social) can mediate this relationship. Focusing in on the peri-trauma phase will help improve understanding of how parents' own appraisals of the child's medical event and coping assistance may influence how children appraise and cope during the medical event. By improving understanding of these factors, preventive interventions can be better tailored and thereby more effective.

Acknowledgements

This work was supported by a Mentored Career Award grant 1K23MH093618-01A1 from the National Institute of Mental Health. Additionally, we would like to thank Aimee Hildenbrand and Katherine Donlon for supporting this manuscript preparation.

References

- Alisic E, Conroy R, Magyar J, Babl FE, O'Donnell ML. Psychosocial care for seriously injured children and their families: A qualitative study among Emergency Department nurses and physicians. *Injury*. (0). doi: <http://dx.doi.org/10.1016/j.injury.2014.02.015>.
- Alisic E, Jongmans MJ, van Wesel F, Kleber RJ. Building child trauma theory from longitudinal studies: a meta-analysis. *Clin Psychol Rev*. 2011; 31(5):736–747. doi: S0272-7358(11)00049-3 [pii]10.1016/j.cpr.2011.03.001. [PubMed: 21501581]
- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, 4th Edition (DSM-IV). Washington, DC: 1994.
- American Psychiatric Association. DSM-5 Development: 309.81 Posttraumatic Stress Disorder. 2010. from <http://www.dsm5.org/ProposedRevisions/Pages/proposedrevision.aspx?rid=165>
- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, 5th Edition (DSM-5). Washington, DC: 2014.
- Balluffi A, Kassam-Adams N, Kazak A, Tucker M, Dominguez T, Helfaer M. Traumatic stress in parents of children admitted to the pediatric intensive care unit. *Pediatric Critical Care Medicine*. 2004; 5:547–553. [PubMed: 15530191]
- Barrett P, Dadds M, Rapee R. Family treatment of childhood anxiety: A controlled trial. *Journal of Consulting and Clinical Psychology*. 1996; 64(2):333–342. [PubMed: 8871418]
- Barrett P, Rapee R, Dadds M, Ryan S. Family enhancement of cognitive style in anxious and aggressive children. *Journal of Abnormal Child Psychology*. 1996; 24(2):187–203. [PubMed: 8743244]
- Berkowitz SJ, Stover CS, Marans SR. The Child and Family Traumatic Stress Intervention: secondary prevention for youth at risk of developing PTSD. *Journal of child psychology and psychiatry, and allied disciplines*. 2011; 52(6):676–685. doi: 10.1111/j.1469-7610.2010.02321.x.
- Blount R, Corbin S, Sturges J, Wolfe V, Prater J, James L. The relationship between adults behavior and child coping and distress during BMA/LP procedures: A sequential analysis. *Behavior Therapy*. 1989; 20:585–601.

- Blount R, Davis N, Powers S, Roberts M. The influence of environmental factors and coping style on children's coping and distress. *Clinical Psychology Review*. 1991; 11:93–116.
- Bryant R, Salmon K, Sinclair E, Davidson P. Heart rate as a predictor of posttraumatic stress disorder in children. *General Hospital Psychiatry*. 2007a; 29:66–68. [PubMed: 17189749]
- Bryant R, Salmon K, Sinclair E, Davidson P. A prospective study of appraisals in childhood posttraumatic stress disorder. *Behaviour Research and Therapy*. 2007b; 45:2502–2507. [PubMed: 17560541]
- Calhoun PS, Hertzberg JS, Kirby AC, Dennis MF, Hair LP, Dedert EA, Beckham JC. The effect of draft DSM-V criteria on posttraumatic stress disorder prevalence. *Depression and anxiety*. 2012; 29(12):1032–1042. Research Support, N.I.H., Extramural. doi: 10.1002/da.22012. [PubMed: 23109002]
- Carrion VG, Weems CF, Ray RD, Glaser B, Hessel D, Reiss AL. Diurnal salivary cortisol in pediatric posttraumatic stress disorder. *Biological Psychiatry*. 2002; 51(7):575–582. doi: [http://dx.doi.org/10.1016/S0006-3223\(01\)01310-5](http://dx.doi.org/10.1016/S0006-3223(01)01310-5). [PubMed: 11950459]
- Cox C, Kenardy J. A randomised controlled trial of a web-based early intervention for children and their parents following accidental injury. *Journal of Pediatric Psychology*. 2010; 35:581–592. doi: 10.1093/jpepsy/jsp095. [PubMed: 19906829]
- Cox C, Kenardy J, Hendrikz J. A meta-analysis of risk factors that predict psychopathology following accidental trauma. *Journal for Specialists in Pediatric Nursing*. 2008; 13(2):98–110. [PubMed: 18366377]
- Dadds M, Barrett P, Rapee R, Ryan S. Family process and child anxiety and aggression: An observational analysis. *Journal of Abnormal Child Psychology*. 1996; 24(6):715–735. [PubMed: 8970906]
- Dalgleish T, Meiser-Stedman R, Kassam-Adams N, Ehlers A, Winston FK, Smith P, Yule W. Predictive validity of acute stress disorder in children and adolescents. *British Journal of Psychiatry*. 2008; 192:392–393. [PubMed: 18450669]
- De Young A, Kenardy J, Spence S. Elevated heart rate as a predictor of PTSD six months following accidental pediatric injury. *Journal of Traumatic Stress*. 2007; 20(5):5.
- Delahanty D, Nugent N, Christopher N, Walsh M. Initial urinary epinephrine and cortisol levels predict acute PTSD symptoms in child trauma victims. *Psychoneuroendocrinology*. 2005; 30:121–128. [PubMed: 15471610]
- Ehlers A, Clark D. A cognitive model of posttraumatic stress disorder. *Behaviour Research and Therapy*. 2000; 38(4):319–345. [PubMed: 10761279]
- Ehlers A, Mayou R, Bryant B. Cognitive predictors of posttraumatic stress disorder in children: results of a prospective longitudinal study. *Behaviour Research and Therapy*. 2003; 41(1):1–10. [PubMed: 12488116]
- Engel GL. The need for a new medical model: a challenge for biomedicine. *Science*. 1977; 196(4286):129–136. [PubMed: 847460]
- Fava GA, Sonino N. The biopsychosocial model thirty years later. *Psychotherapy and psychosomatics*. 2008; 77(1):1–2. Biography Editorial Historical Article. doi: 10.1159/000110052. [PubMed: 18087201]
- Finkelhor D, Turner H, Ormrod R, Hamby SL. Violence, abuse, and crime exposure in a national sample of children and youth. *Pediatrics*. 2009; 124(5):1411–1423. Research Support, U.S. Gov't, Non-P.H.S. doi: 10.1542/peds.2009-0467. [PubMed: 19805459]
- Folkman S, Lazarus R. If It Changes It Must Be a Process: Study of Emotion and Coping During Three Stages of a College Examination. *Journal of Personality and Social Psychology*. 1985; 48(1):150–170. [PubMed: 2980281]
- Gold JI, Kant AJ, Kim SH. The impact of unintentional pediatric trauma: A review of pain, acute stress, and posttraumatic stress. *Journal of Pediatric Nursing*. 2008; 23(2):81–91. doi: <http://dx.doi.org/10.1016/j.pedn.2007.08.005>. [PubMed: 18339334]
- Grossman D. The history of injury control and the epidemiology of child and adolescent injuries. *The Future of Children*. 2000; 10(1):23–52. [PubMed: 10911687]

- Holbrook T, Hoyt D, Coimbra R, Potenza B, Sise M, Anderson J. Long-term posttraumatic stress disorder persists after major trauma in adolescents: New data on risk factors and functional outcome. *Journal of Trauma-Injury Infection and Critical Care*. 2005; 58(4):764–769.
- Kahana S, Feeny N, Youngstrom E, Drotar D. Posttraumatic stress in youth experiencing illnesses and injuries: An exploratory meta-analysis. *Traumatology*. 2006; 12(2):148–161.
- Kassam-Adams N, Garcia-España J, Fein J, Winston FK. Heart rate and posttraumatic stress in injured children. *Archives of General Psychiatry*. 2005; 62:335–340. [PubMed: 15753247]
- Kassam-Adams N, Marsac M, Hildenbrand A, Winston FK. Posttraumatic stress following pediatric injury: Update on diagnosis, risk factors, and intervention. *JAMA Pediatrics*. 2013; 167(12):1158–1165. doi: 10.1001/jamapediatrics.2013.2741. [PubMed: 24100470]
- Kassam-Adams N, Marsac ML, Cirilli C. PTSD symptom structure in injured children: Relationships with functional impairment and depression symptoms. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2010; 49(6):616–625. [PubMed: 20494271]
- Kassam-Adams N, Winston FK. Predicting child PTSD: The relationship between acute stress disorder and PTSD in injured children. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2004; 43(4):403–411. [PubMed: 15187800]
- Kazak A. Pediatric Psychosocial Preventative Health Model (PPPHM): Research, practice, and collaboration in pediatric family systems medicine. *Families, Systems, & Health*. 2006; 24(4):381–395.
- Kazak A, Kassam-Adams N, Schneider S, Zelikovsky N, Alderfer M, Rourke M. An integrative model of pediatric medical traumatic stress. *Journal of Pediatric Psychology*. 2006; 44:343–355. [PubMed: 16093522]
- Kean E, Kelsay K, Wamboldt F, Wamboldt M. Posttraumatic stress in adolescents with asthma and their parents. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2006; 45:78–86. [PubMed: 16327584]
- Klaassens ER, Giltay EJ, Cuijpers P, van Veen T, Zitman FG. Adulthood trauma and HPA-axis functioning in healthy subjects and PTSD patients: a meta-analysis. *Psychoneuroendocrinology*. 2012; 37(3):317–331. Meta-Analysis Research Support, Non-U.S. Gov't. doi: 10.1016/j.psyneuen.2011.07.003. [PubMed: 21802212]
- Kliewer W, Parrish K, Taylor K, Jackson K, Walker J, Shivy V. Socialization of coping with community violence: Influences of caregiver coaching, modeling, and family context. *Child Development*. 2006; 77(3):605–523. [PubMed: 16686791]
- Kramer DN, Landolt MA. Characteristics and efficacy of early psychological interventions in children and adolescents after single trauma: a meta-analysis. *European Journal of Psychotraumatology*. 2011; 2:24. doi: 10.3402/ejpt.v2i0.7858.
- Landolt M, Vollrath M, Gnehm H, Sennhauser F. Post-traumatic stress impacts on quality of life in children after road traffic accidents: Prospective study. *Australian and New Zealand Journal of Psychiatry*. 2009; 43(8):746–753. [PubMed: 19629796]
- Landolt M, Vollrath M, Ribi K. Predictors of coping strategy selection in paediatric patients. *Acta paediatr*. 2002; 91:945–960.
- Lazarus, R. *Emotion and adaptation*. Oxford University Press; New York: 1991.
- Marks, JS.; McQueen, DV. Chronic disease. In: Koop, CE.; Pearson, CE.; Schwarz, MR., editors. *Critical Issues in Global Health*. Josey-Bass; San Francisco: 2001. p. 117-126.
- Marsac ML, Donlon KA, Hildenbrand AK, Winston FK, Kassam-Adams N. Understanding recovery in children following traffic-related injuries: Exploring acute traumatic stress reactions, child coping, and coping assistance. *Clinical Child Psychology and Psychiatry*. 2014; 19(2):233–243. doi: 10.1177/1359104513487000. [PubMed: 23677925]
- Marsac ML, Donlon KA, Winston FK, Kassam-Adams N. Child coping, parent coping assistance, and post-traumatic stress following paediatric physical injury. *Child: Care, Health and Development*. 2013; 39(2):171–177. doi: 10.1111/j.1365-2214.2011.01328.x.
- McDermott B, Cvitanovich A. Posttraumatic stress disorder and emotional problems in children following motor vehicle accidents: An extended case series. *Australian and New Zealand Journal of Psychiatry*. 2000; 34:446–452. [PubMed: 10881968]

- McNally R, Bryant R, Ehlers A. Does early psychological intervention promote recovery from posttraumatic stress? *Psychological Science in the Public Interest*. 2003; 4(2):45–79.
- Meewisse ML, Reitsma JB, de Vries GJ, Gersons BP, Olff M. Cortisol and post-traumatic stress disorder in adults: systematic review and meta-analysis. *The British journal of psychiatry : the journal of mental science*. 2007; 191:387–392. Meta-Analysis Review. doi: 10.1192/bjp.bp.106.024877. [PubMed: 17978317]
- Meiser-Stedman R. Towards a cognitive -behavioral model of PTSD in children and adolescents. *Clinical Child and Family Psychology*. 2002; 5(4):217–232.
- Meiser-Stedman R, Dalgleish T, Glucksman E, Yule W, Smith P. Maladaptive cognitive appraisals mediate the evolution of posttraumatic stress reactions: A 6-month follow-up of child and adolescent assault and motor vehicle accident survivors. *Journal of Abnormal Psychology*. 2009; 118(4):9.
- Meiser-Stedman R, Smith P, Yule W, Dalgleish T, Glucksman E. Diagnostic, demographic, memory quality, and cognitive variables associated with acute stress disorder in children and adolescents. *Journal of Abnormal Psychology*. 2007; 116:65–79. [PubMed: 17324017]
- Micco J, Ehrenreich J. Children's interpretation and avoidant response biases in response to non-salient and salient situations: Relationships with mothers' threat perception and coping expectations. *Journal of Anxiety Disorders*. 2008; 22(3):371–385. doi: DOI: 10.1016/j.janxdis.2007.03.009. [PubMed: 17434288]
- Miller, MW.; Wolfe, EJ.; Fabricant, L.; Stein, N. Low Basal Cortisol and Startle Responding as Possible Biomarkers of PTSD: The Influence of Internalizing and Externalizing Comorbidity. In: Shiromani, P.; Keane, T.; LeDoux, JE., editors. *Post-Traumatic Stress Disorder: Basic Science and Clinical Practice*. Humana Press; 2009. p. 277–293.
- Mintzer L, Stuber M, Seacord D, Castaneda M, Mesrkhani V, Glover D. Traumatic stress symptoms in adolescent organ transplant recipients. *Pediatrics*. 2005; 115(6):1640–1649. [PubMed: 15930227]
- Murray, C.; Lopez, A. *The Global Burden of Disease: A Comprehensive Assessment of Mortality and Disability from Diseases, Injuries, and Risk Factors in 1990 and Projected to 2020*. Harvard University Press; Cambridge, Massachusetts: 1996.
- Nixon RDV, Nehmy TJ, Ellis AA, Ball S-A, Menne A, McKinnon AC. Predictors of posttraumatic stress in children following injury: The influence of appraisals, heart rate, and morphine use. *Behaviour Research and Therapy*. 2010; 48(8):810–815. doi: http://dx.doi.org/10.1016/j.brat.2010.05.002. [PubMed: 20537316]
- Nugent N, Christopher N, Delahanty D. Emergency medical service and in-hospital vital signs as predictors of subsequent PTSD symptom severity in pediatric injury patients. *Journal of Child Psychology and Psychiatry*. 2006a; 47:919–926. [PubMed: 16930386]
- Nugent N, Christopher N, Delahanty D. Initial physiological responses and perceived hyperarousal predicts subsequent emotional numbing in child trauma victims. *Journal of Traumatic Stress*. 2006b; 19:349–359. [PubMed: 16789001]
- Olff M, Langeland W, Gersons B. The psychology of PTSD: Coping with trauma. *Psychoneuroendocrinology*. 2005; 30:8.
- Ostrowski M, Christopher N, van Dulmen M, Delahanty D. Acute child and mother psychophysiological responses and subsequent PTSD following a child's traumatic event. *Journal of Traumatic Stress*. 2007; 20:1–11.
- Ostrowski SA, Ciesla JA, Lee TJ, Irish L, Christopher NC, Delahanty DL. The impact of caregiver distress on the longitudinal development of child acute post-traumatic stress disorder symptoms in pediatric injury victims. *Journal of Pediatric Psychology*. 2011; 36(7):806–815. Research Support, N.I.H., Extramural. doi: 10.1093/jpepsy/jsq113. [PubMed: 21262744]
- Pitman RK, Delahanty DL. Conceptually driven pharmacologic approaches to acute trauma. *CNS spectrums*. 2005; 10(2):99–106. Research Support, N.I.H., Extramural Research Support, U.S. Gov't, P.H.S. [PubMed: 15685120]
- Pitman RK, Rasmusson AM, Koenen KC, Shin LM, Orr SP, Gilbertson MW, Liberzon I. Biological studies of post-traumatic stress disorder. *Nature reviews. Neuroscience*. 2012; 13(11):769–787. Review. doi: 10.1038/nrn3339.

- Polinder S, Meerdering WJ, Toet H, Mulder S, Essink-Bot ML, van Beeck EF. Prevalence and prognostic factors of disability after childhood injury. *Pediatrics*. 2005; 116(6):e810–817. doi: 10.1542/peds.2005-1035. [PubMed: 16322139]
- Power T. Stress and coping in childhood: The parents' role. *Parenting: Science and Practice*. 2004; 4(4):271–317.
- Rose S, Bisson J, Churchill R, Wessely S. Psychological debriefing for preventing post traumatic stress disorder (PTSD). *Cochrane Database Syst Rev*. 2002; (2) CD000560. doi: CD000560 [pii] 10.1002/14651858.CD000560.
- Salmon K, Bryant R. Posttraumatic stress disorder in children: The influence of developmental factors. *Clinical Psychology Review*. 2002; 22(2):163–188. doi: DOI: 10.1016/S0272-7358(01)00086-1. [PubMed: 11806018]
- Schwartz E, Perry B. The post-traumatic response in children and adolescents. *Psychiatric Clinics of North America*. 1994; 17(2):311–326. [PubMed: 7937362]
- Skinner E, Zimmer-Gembeck M. The development of coping. *Annual Review of Psychology*. 2007; 58:119–144.
- Stallard P, Smith E. Appraisals and cognitive coping styles associated with chronic post-traumatic symptoms in child road traffic accident survivors. *Journal of child psychology and psychiatry, and allied disciplines*. 2007; 48(2):194–201. doi: JCPP1692 [pii] 10.1111/j.1469-7610.2006.01692.x.
- Stallard P, Velleman R, Salter E, Howse I, Yule W, Taylor G. A randomised controlled trial to determine the effectiveness of an early psychological intervention with children involved in road traffic accidents. *Journal of Child Psychology and Psychiatry*. 2006; 47(2):127–134. [PubMed: 16423143]
- Stoddard F, Saxe G. Ten year research review of physical injuries. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2001; 40(10):1128–1145. [PubMed: 11589526]
- Suls J, Rothman A. Evolution of the biopsychosocial model: prospects and challenges for health psychology. *Health psychology : official journal of the Division of Health Psychology, American Psychological Association*. 2004; 23(2):119–125. Research Support, U.S. Gov't, Non-P.H.S. Research Support, U.S. Gov't, P.H.S. doi: 10.1037/0278-6133.23.2.119.
- Trickey D, Siddaway AP, Meiser-Stedman R, Serpell L, Field AP. A meta-analysis of risk factors for post-traumatic stress disorder in children and adolescents. *Clinical Psychology Review*. 2012; 32(2):122–138. Meta-Analysis Research Support, Non-U.S. Gov't. doi: 10.1016/j.cpr.2011.12.001. [PubMed: 22245560]
- Walker A, Harris G, Baker A, Kelly D, Houghton J. Posttraumatic stress responses following liver transplantation in older children. *Journal of Child Psychology and Psychiatry*. 1999; 40(3):363–374. [PubMed: 10190338]
- Weems CF, Carrion VG. The association between PTSD symptoms and salivary cortisol in youth: the role of time since the trauma. *Journal of Traumatic Stress*. 2007; 20(5):903–907. doi: 10.1002/jts.20251. [PubMed: 17955527]
- Zatzick D, Jurkovich G, Fan M, Grossman D, Russo J, Katon W, Rivara F. Association between posttraumatic stress and depressive symptoms and functional outcomes in adolescents followed up longitudinally after injury hospitalization. *Archives of Pediatrics & Adolescent Medicine*. 2008; 162(7):642–648. [PubMed: 18606935]
- Zehnder D, Meuli M, Landolt M. Effectiveness of a single-session early psychological intervention for children after road traffic accidents: A randomised controlled trial. *Child and Adolescent Psychiatry and Mental Health*. 2010; 4:7. [PubMed: 20181120]
- Zehnder D, Prchal A, Vollrath M, Landolt M. Prospective study of the effectiveness of coping in pediatric patients. *Child Psychiatry and Human Development*. 2006; 36(3):351–368. [PubMed: 16362238]
- Zonfrillo MR, Durbin DR, Winston FK, Zhao H, Stineman MG. Physical disability after injury-related inpatient rehabilitation in children. *Pediatrics*. 2013; 131(1):e206–213. Multicenter Study Research Support, Non-U.S. Gov't. doi: 10.1542/peds.2012-1418. [PubMed: 23248228]




Figure 1. Theoretical model for the development of child PTSD/PTSS following acute medical trauma*

* This model does not include every factor influencing the development of PTSS following acute medical trauma. Rather, we suggest including pre, peri, and post-trauma factors in future research and highlight specific understudied paths during the peri-trauma period. See text for supporting evidence for individual paths as labeled by letters (a-o).

Table 1

Summary of meta-analyses examining predictors of PTSD/PTSS in children

	Kahana et al. (2006)	Cox et al. (2008)	Alisic et al. (2011)	Trickey et al. (2011)
Meta-analysis Characteristics				
Traumatic Events	Injury & illness	Injury	Various	Various
Number of Studies	26	14	34	62
Designs of Studies	Cross-sectional	Cross-sectional & longitudinal	Longitudinal	Cross-sectional
Age in Samples	6-19	5-18	1-18	6-18
PTSD Measures	Included all PTSD symptoms	Validated/ psychometrics provided	Included at least avoidance and intrusion	Validated/ include intrusion, avoidance, hyperarousal
Model	Fixed effects	Fixed effects	Random effects	Random effects
Predictors Assessed	Effect Sizes			
Pre-Trauma Factors				
<i>Biological</i>				
Sex (female)	0	Small	Small	Small
Age (younger)	0 – Medium	0 – Small	0	0
Minority status	--	Small	0	0
<i>Psychological</i>				
Pre-trauma psychopathology	--	Small	--	Small
Pre-trauma low self esteem	--	--	--	Small
<i>Social (Environmental)</i>				
Low SES	Small	--	0	Small
Prior trauma exposure	--	Small	--	--
Other life events	--	--	--	Small
Parent psychological problem	--	--	--	Small
Peri-Trauma Factors				
<i>Biological</i>				
Heart rate at hospital	--	--	Small	--
<i>Psychological</i>				
Perceived life threat	Large	Small- Medium	--	Small
Peri-trauma fear	--	--	--	Small
Appraisal	Small	--	--	--
<i>Social (Environmental)</i>				
Trauma severity	Small – Large	0 – Small	0	Small
Time since trauma (shorter)	--	--	--	Small
Days in hospital	--	--	Small	--
Involvement of family/ friend in trauma		0 – Small		

	Kahana et al. (2006)	Cox et al. (2008)	Alisic et al. (2011)	Trickey et al. (2011)
Bereavement	--	--	--	Small
Post-trauma Factors				
<i>Biological</i>				
No factors addressed	--	--	--	--
<i>Psychological</i>				
Co-morbid psych problem	--	--	--	Small
Acute stress	Large	--	Medium	Medium
Depressive symptoms	Large	--	Medium	Small – Medium
Anxiety symptoms	Large	--	Medium	Small – Medium
Dissociation	Large	--	--	--
Internalizing Symptoms	Large	--	--	--
Conduct problems	--	--	--	--
Coping: Blaming Others	--	--	--	Medium
Coping: Distraction	--	--	--	Medium
Coping: Thought Suppression				Medium
<i>Social (Environmental)</i>				
Parental distress	--	0 – Medium	Small	Small
Poor family functioning	--	--	--	Medium
Social Impairment	Small	--	--	--
Social Support (low)	Small	--	--	Small
Social Withdrawal	--	--	--	Small

Table 2

Support for potential peri-trauma phase predictors of subsequent PTSS/PTSD

Predictors	Supporting Study Citations
<i>Biological</i>	
Memory formation	Pitman & Delahanty, 2005 ^a ; Pitman et al., 2012 ^a
Elevated cortisol (boys only)	Carrion et al., 2002; Delahanty, et al., 2005; Ostrowski, et al., 2007
Elevated heart rate	Nugent et al., 2006 ^a ; Kassam-Adams, et al., 2005; Bryant et al., 2007; De Young et al., 2007; Nugent et al., 2006 ^b
<i>Psychological</i>	
Child PTSS	Ostrowski et al., 2011
Child cognitive appraisals	Kahana et al., 2006; Cox et al., 2008; Trickey et al., 2012; Ehlers et al., 2003; McDermott & Cvitanovich, 2000; Stallard & Smith, 2007; Bryant et al., 2007; Aaron et al., 1999
Child coping	Stallard & Smith, 2007; Zehnder et al., 2006; Marsac et al., 2013; Marsac et al., 2014; Landolt et al., 2002
<i>Social (Environmental)</i>	
Trauma severity	Kahana et al., 2006; Cox et al., 2008; Trickey et al., 2012
Medical Team Support	Kazak, 2006 ^b ; Kassam-Adams et al., 2013 ^b
Community Support	Kazak, 2006 ^b ; Kassam-Adams et al., 2013 ^b
Parent PTSS	Balluffi et al., 2004; Alisic et al., 2011; Cox et al., 2008; Trickey et al., 2012
Parent appraisals	Barrett et al., 1996 ^c ; Dadds et al., 1996 ^c
Parent coping assistance	Marsac et al., 2013; Barrett et al., 1996 ^c ; Dadds et al., 1996 ^c

^a Studies conducted during post-trauma period only; based on theory, we expect findings to be similar during peri-trauma;

^b Suggested by theory; no empirical evidence to date;

^c Studies were conducted to study how appraisals and coping were influenced by parents in relation to anxiety as an outcome. While samples were not children presenting with trauma-exposure, we suggest that results specific to how parents influence child appraisals and coping will apply to families in which children have been exposed to medical trauma.