

Becoming Trauma Informed: Validating a Tool to Assess Health Professional's Knowledge, Attitude, and Practice

Simmy King, DNP, MS, MBA, RN-BC, NE-BC*; Kuan-Lung Daniel Chen, DrPH(c), MPH†; Binny Chokshi, MD‡§

Abstract

Introduction: To effectively address the negative health effects of early childhood trauma and adversity, healthcare professionals and healthcare institutions must understand the impact of adverse childhood experiences and trauma on health. This study aimed to validate a tool to assess knowledge, attitude, and practice of trauma-informed care among interdisciplinary pediatric healthcare staff.

Methods: A 36-item survey tool, "Knowledge, Attitudes, and Practices of Trauma-Informed Practice," was adopted and modified with permission from the author. We administered the survey electronically to 2,659 staff at a pediatric healthcare institution. To assess the tool's reliability and validity, internal consistency reliability testing, content validity, and construct validity assessments were conducted. **Results:** A total of 592 surveys were collected, representing a 22.3% response rate. Confirmatory factor analysis revealed that 21 items provided the strongest internal consistency reliability for the overall tool and each factor. The overall Cronbach's alpha for the 21-item tool was 0.86, with 0.84 for the knowledge factor, 0.74 for the attitude factor, and 0.78 for the practice factor. The goodness of fit based on this analysis was good to adequate, with a 0.077 root mean square error of approximation. **Conclusions:** Healthcare professionals and organizations are in a unique position to improve the health and well-being of their patients by implementing a trauma-informed approach to minimize the impact of adverse childhood experiences and trauma. This validated tool will allow organizations to identify gaps in knowledge, attitude, and practice among staff to subsequently begin developing pointed strategies to achieve a culture of trauma-informed practice. (*Pediatr Qual Saf* 2019;4:e215; doi: 10.1097/pq9.000000000000215; Published online September, 9 2019.)

INTRODUCTION

In the United States, two-thirds of individuals have been exposed to ≥ 1 traumatic event in their childhood. These exposures to trauma can place children at risk for emotional, physical, and functional impairment.¹ Although there are many definitions of trauma, the most commonly



From the *Department of Nursing, Children's National Health System, Washington, D.C.; †Sumner M. Redstone Global Center for Prevention and Wellness, Milken Institute School of Public Health, The George Washington University, Washington, D.C.; ‡Goldberg Center for Community Pediatric Health, Division of Adolescent and Young Adult Medicine, Children's National Health System, Washington, D.C.; and §Department of Pediatrics, George Washington University School of Medicine and Health Sciences, Washington, D.C.

*Corresponding author. Address: Simmy King, DNP, MS, MBA, RN-BC, NE-BC, Department of Nursing, Children's National Health System, 111 Michigan Avenue NW, Washington, DC 20010 Email: simmy.king@childrensnational.org PH: 202-476-4151; fax: 202-476-5935

Copyright © 2019 the Author(s). Published by Wolters Kluwer Health, Inc. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

To cite: King S, Chen KLD, Chokshi B. Becoming Trauma Informed: Validating a Tool to Assess Health Professional's Knowledge, Attitude, and Practice. *Pediatr Qual Saf* 2019;5:e215.

Received for publication March 5, 2019; Accepted August 15, 2019.

Published online September 9, 2019

DOI: 10.1097/pq9.000000000000215

referred to definition is from the Substance Abuse and Mental Health Services Administration. They define trauma as "an event, series of events, or set of circumstances experienced by an individual as physically or emotionally harmful or life-threatening with lasting adverse effects on the individuals functioning and mental, physical, social, emotional, or spiritual well being."² Although this definition provides context for healthcare professionals, it is important to note that trauma is a subjective experience; how an individual responds to trauma may evolve and change over time and well into adulthood.³ The seminal study conducted by Felitti et al⁴ underscores the impact of 10 specific childhood traumas, now commonly known as "adverse childhood experiences" (ACEs), by demonstrating how exposure to ACEs can lead to risky health behaviors and poor health outcomes in a dose-dependent manner. These findings, together with current research regarding the negative impact of childhood experiences on the future health and functioning of pediatric patients, serve as a call to action for healthcare professionals to incorporate trauma-informed care (TIC) into their routine clinical practice.⁴⁻⁷

TIC is an approach to healthcare that is rooted in the realization of the widespread incidence of trauma, recognition of the signs and symptoms of trauma, response

to trauma in a full and integrated manner, and resisting retraumatization by organizations and healthcare professionals.² Furthermore, TIC includes adherence to the following key principles: (1) safety, (2) trustworthiness, (3) peer support, (4) collaboration, (5) empowerment and choice, and (6) cultural, historical, and sex issues.^{2,8} TIC has historically been used among health professionals in social work, mental health, child services, and the juvenile justice program to inform their work, but it has not been routinely practiced by pediatric healthcare professionals.^{1,8} Now with increasing evidence that TIC can lead to improved patient care outcomes, healthcare institutions are recognizing the importance of clinician competency in TIC.^{9,10} In 2014, the AAP's (American Academy of Pediatrics) "Trauma Toolbox for Primary Care" highlighted the role of a trauma-informed approach in addressing childhood adversity and trauma in the pediatric primary care setting.¹¹ Subsequent literature has highlighted the need for pediatric healthcare organizations to advance TIC¹² by educating healthcare professionals and developing competency in TIC.¹³

Regional data from the 2016 National Child Health Survey showed that 47% of the children in our region have had ≥ 1 ACE.¹⁴ These data underscored the need for a coordinated approach to address trauma in the pediatric patients we care for at Children's National Health System. As a first step, we assessed the learning needs of our target audience, a crucial first step in developing a learning curriculum.¹⁵ Specifically, we wanted to measure knowledge, attitude, and practice (KAP) as they related to TIC to inform our educational intervention. The KAP method is a reliable and valid method to enhance KAP around a specific theme, establish baseline data, and suggest learner-centric intervention strategies.¹⁶ For larger organizations, the best method of conducting a formal learning needs gap analysis is to use a valid and reliable survey tool.¹⁵ Therefore, we searched for a validated tool for our needs assessment.

From our literature review, we found one study that measured nursing knowledge and experience related to TIC, but it was a qualitative study.¹⁷ Another study by Bruce et al¹⁸ surveyed providers working at a trauma center regarding TIC; however, this survey included questions related to posttraumatic stress, and these questions were not aligned with the intent of our work. Baker et al¹⁹ was the only article that developed and validated a tool to measure TIC, the ARTIC (Attitudes Related to Trauma-Informed Care Scale). In this study, psychometric analysis established strong internal consistency and test-retest reliability for 45-item form with 7 subscales and a 35-item composite and 10-item short form. Although this tool was intended for use across disciplines, participants in the original study worked in education and human service settings.

Additionally, survey items were geared to measure the extent to which an individual or system is trauma-informed, with a lack of focus on understanding the important link between traumatic exposures and health outcomes. Missing from the available literature was a

validated tool that measured KAP related to TIC among healthcare professionals within a healthcare setting. As such, this study aimed to validate a tool to assess KAP of TIC among interdisciplinary pediatric healthcare staff.

METHODS

Adopting Existing Survey Instrument

After a comprehensive literature review, we identified a survey tool that we could adapt for widespread use among healthcare professionals. The "Knowledge, Attitudes, and Practices of Trauma-Informed Practice Survey" is a 36-item unpublished and unvalidated tool developed by Abdoh et al.²⁰ Abdoh et al²⁰ designed the tool based on a literature review, then conducted face validity assessment with staff at their community center and pilot-tested the tool with 17 multidisciplinary staff members at that site. The community center provides housing to individuals with substance dependencies and concurrent physical and mental health issues. Its staff consist of various professional disciplines, including physicians, social workers, nurses, and mental health support staff. Among the 36 survey items, 29 questions measure knowledge, attitudes, and practice; 12 items related to knowledge, 11 related to attitudes, and 6 related to practices. Each item has a 5-point Likert scale response format with options from *strongly disagree* to *strongly agree*. The other 7 items of the survey capture demographic data, including the respondent's role, department, years of experience, age group, and previous experience with TIC. We adopted this survey for our learning needs assessment after discussion with, and permission from, the author.

Face and Content Validity

To establish face validity, internal experts in the field of childhood adversity and trauma reviewed each item to assess the degree to which it would measure each of the 3 factors: KAP related to TIC. Internal expert qualifications included serving as faculty facilitators for continuing medical education on TIC institutionally, peer-reviewed publications related to mental health and trauma, employing TIC principles in their pediatric practice, coordinating federal and private funding on childhood trauma, and presenting at national conferences. We provided the following definitions of knowledge, attitudes, and practice to the internal experts as reference. Knowledge was defined as the degree to which staff members were aware of, informed about, or familiar with the applicability of TIC in the healthcare setting. Attitude was defined as how staff felt or thought about TIC, and practice was defined as how staff applied their knowledge and attitudes about TIC into their practice. We also asked these experts to review and assess whether the items on the tool should be included. Based on the review by the internal experts, only 1 item in the original tool, "Trauma-informed practice (TIP) shares many similarities to harm reduction," was removed from the survey because our survey was being implemented in a pediatric healthcare institution whose

staff is not well versed in harm reduction techniques. Additionally, the word “client” was replaced with “patient” to better align with the context of a large health-care institution, versus a community setting. When determining content validity, institutional experts considered the item’s (1) relevance to TIP in general, (2) relevance to its associated factor, (3) alignment with common wording used at the institution, and (4) clarity of wording.

Data Collection

Before data collection, the organizational Institutional Review Board reviewed and approved the study. We collected the initial data between November and December 2017. An online survey platform, SurveyMonkey® (San Mateo, Calif.), was used to administer the survey tool, and the initial request to participate was sent via hospital email, followed by 2 subsequent email reminders until the survey closing date.

Construct Validity: Confirmatory Factor Analysis

The survey tool was adopted specifically for its ability to assess the 3 latent factors: knowledge, attitudes, and practices. Therefore, the priority for tool validation was to investigate whether the survey items are indeed manifest variables of the 3 factors.

Using confirmatory factor analysis (CFA) to assess the association between survey items and the 3 theorized latent factors, we assigned numeric values to each response for the analysis: *Strongly disagree* was assigned the value of 0, *disagree* a value of 1, *neutral* a value of 2, *agree* a value of 3, and *strongly agree* a value of 4. We assessed model fit based on factor loadings for individual survey items and goodness of fit tests commonly used in literature,^{21–23} including chi-square test, the Bentler comparative fit index^{24,25} (CFI), Tucker–Lewis index (TLI), and root mean square error of approximation (RMSEA). Akaike information criterion and Bayesian information criterion were also used to compare models. We tested survey items with a high residual for association with different factors based on modification indices.²⁶ Items sharing a standard covariance residual of ≥ 1.96 would prompt a review of the modification indices. If the item’s modification index was >3.84 , it would then prompt an attempt to assess whether the item fits better with a different factor.²⁷ We removed items if the model fit did not improve as a result of changing the associated factor.^{26,27}

Reliability: Internal Consistency

We assessed internal consistency by calculating Cronbach’s alpha for each factor and the overall tool each time CFA produced an acceptable model.²⁸ Using a combination of CFA and Cronbach’s alpha for survey tool validation is consistent with similar research in the field.^{23,28}

RESULTS

Survey Respondents: Descriptive Data

We distributed the survey to 2,659 staff members. A total of 592 survey responses were collected—representing a

22.3% response rate. Among the responses collected, 37 were removed for no response on scale items, and 4 were removed for repeat entries. We deleted entries with missing scale item data listwise. Among the remaining 511 respondents, 54.4% were nurses, and 19.2% were attending physicians. Age distribution among the respondents was mostly even. A quarter of respondents (24.1%) had <2 years of experience in their current role, and 41.7% had ≥ 10 years in their current role. Approximately one-third of the respondents received some TIP training before completing the survey (Table 1).

Factor Analysis

We conducted CFA using Stata 15 software (StataCorp LLC, College Station, TX) to assess model fit and adjust the model to improve fit. The sample exceeds the commonly recommended minimum size of 300.²⁹ Data distribution for all items, except for one, meet multivariate normality. Therefore, we use maximum likelihood, which allows for mild departure for multivariate normality, as the estimation method for CFA.²¹ As part of the model fitting process, we removed 7 items from the survey. The final 21-item model showed adequate overall model fit with a 0.077 RMSEA.²⁸ The chi-square statistics is 748.05 with a P value of <0.001 . However, for sample sizes >500 , it is common for the chi-square test to find statistical significance.²⁹ The CFI is 0.857, and the TLI is 0.839, each falls slightly below the 0.9 cutoff for “good fit.”^{23–25} Altogether,

Table 1. Descriptive Data for the 511 Respondents Whose Questionnaires Were Used for Confirmatory Factor Analysis

Variable	N (%)
Age, y	
20–25	39 (7.63)
26–30	67 (13.11)
31–35	83 (16.24)
36–40	70 (13.70)
41–50	92 (18.00)
50+	160 (31.31)
Role	
Nurse	278 (54.40)
Attending physician	98 (19.18)
Advanced practice nurse	25 (4.89)
Nonphysician other	25 (4.89)
Social worker	16 (3.13)
Psychologist	14 (2.74)
Child psychiatric specialist	12 (2.35)
Emergency/trauma technician	12 (2.35)
Resident physician	12 (2.35)
Fellow physician	9 (1.76)
Psychiatry technician	4 (0.78)
Patient care technician	3 (0.59)
Unit clerk/front desk support	3 (0.59)
Years of experience in current role	
<2	123 (24.07)
2–5	94 (18.40)
6–10	81 (15.85)
>10	213 (41.68)
Participated in any trauma-informed practice training/education since working at Children’s National	
Yes	179 (35.03)
No	331 (64.77)
No response	1 (0.20)

Table 2. Final 21-item Scale to Assess Staff Knowledge, Attitude, and Practice Related to Trauma-informed Care, Following Confirmatory Factor Analysis

Factor	Survey Question	Initial Number*	Factor Loading (Standardized)	Standard Error	Cronbach's alpha
Knowledge	1. Exposure to trauma is common.	8	0.4188	0.0389	0.8592
	2. Trauma affects physical, emotional, and mental well-being.	9	0.6266	0.0295	
	3. Substance use issues can be indicative of past traumatic experiences or ACEs.	11	0.8753	0.0145	
	4. There is a connection between mental health issues and past traumatic experiences or ACEs.	12	0.8218	0.0177	
	5. Distrusting behavior can be indicative of past traumatic experiences or ACEs.	13	0.8247	0.0175	
Attitude	6. Retraumatization can occur unintentionally.	18	0.6264	0.0296	0.7593
	7. Recovery from trauma is possible.	20	0.4750	0.0391	
	8. Paths to healing/recovery from trauma are different for everyone.	21	0.6344	0.0317	
	9. People are experts in their own healing/recovery from trauma.	22	0.3703	0.0424	
	10. Informed choice is essential in healing/recovery from trauma.	23	0.6746	0.0294	
	11. TIP is essential for working effectively with our patients and their families.	25	0.7472	0.0254	
	12. I have a comprehensive understanding of TIP.	26	0.3327	0.0447	
	13. I believe in and support the principles of TIP.	27	0.6691	0.0304	
	14. I share my expertise and collaborate effectively with colleagues regarding the use of TIP.	28	0.4392	0.0411	
	Practice	15. I would like to receive more training on TIP.	30	0.3654	
16. I maintain transparency in all interactions with patients.		31	0.5771	0.0337	
17. I offer patients' choices and respect their decisions.		32	0.7443	0.0249	
18. I help patients and peers to recognize their own strengths.		33	0.7301	0.0257	
19. I inform all patients of my actions before I perform them.		34	0.6992	0.0274	
20. My interaction with each patient is unique and tailored to their specific needs.		35	0.7608	0.0241	
21. I practice self-care (taking care of my own needs and well-being).		36	0.2828	0.0445	
Goodness of fit					
Chi-square	748.05; $P < 0.001$				
RMSEA	0.077				
CFI	0.857				
TLI	0.839				
SRMR	0.067				
CD	0.995				

*Original numbers represent those of Abdoh et al²⁰ and begin with 8 since the first 7 items are demographic questions. Eight items from the original 36-item survey were removed: 10, trauma can have lifelong effects that may span generations; 14, TIP requires providers to recognize, understand, and respond to the effects of trauma; 15, TIP aims to create safe environments that promote healing and recovery from trauma exposure; 16, TIP includes understanding the physical, psychological, and emotional safety of both the patient and the provider; 17, when using TIP, you must know specific details of a patient's history of trauma; 19, retraumatization can occur in both community and institutional settings; 24, TIP shares many similarities to harm reduction (removed after content validity review, before confirmatory factor analysis); and 29, I have all the resources I need to engage in TIP.

ACE, adverse childhood experience; CD, coefficient of determination; SRMR, standardized root mean square residual; TIP, trauma-informed practice.

the indices suggest an acceptable model. Internal consistency was good: Cronbach's alpha values were within acceptable range with values of ≥ 0.74 for each of the 3 factors.³⁰ Specifically, the values were 0.84 for knowledge (with an item-to-total correlation range of 0.42–0.59), 0.74 for attitude (0.37–0.66), and 0.78 for practice (0.30–0.59).

The standardized interfactor correlation coefficients were 0.55 for knowledge–attitude ($P < 0.001$), 0.28 for knowledge–practice ($P < 0.001$), and 0.65 for attitude–practice ($P < 0.001$). The interfactor correlations suggest, by theory, some level of association among KAP factors for TIP. The medium to low value of the interfactor correlations also suggests that it is not redundant for the tool to include all 3 factors.

We reduced the total number of items from 28 to 21 in the final model (Table 2). In the final 21-item model, 7 items measure knowledge (5 items removed), 8 items measure attitude (2 items removed), and 6 items measure practice (0 items removed). The 7 removed items and any subset of the 7 do not form a fourth factor that would be theoretically meaningful.

DISCUSSION

This study aimed to establish the validity of the “Knowledge, Attitude, and Practice Related to Trauma-Informed Practice” tool. Our analysis indicated that the 21-item version could reliably assess KAP related to TIC among healthcare professionals in a pediatric institution. With increased awareness among healthcare professionals regarding the health impact of ACEs and trauma, healthcare organizations will need to effectively assess the learning needs of their staff to address gaps in KAP to implement a TIC approach that meets the needs of their patients.

At our institution, we used our survey results to develop and deliver an educational intervention for hospital staff. We partnered with the National Center on Trauma-Informed Care and Alternatives to Restraint and Seclusion, part of the Substance Abuse and Mental Health Services Administration, to provide 7 half-day learning sessions between January and March 2018. We have trained over 700 staff members and measured pre and post KAP using the validated KAP related to TIC tool. Our next steps include analyzing postintervention

data and performing additional tool analysis that we will share in a future article.

Limitations

The first limitation of this work is that we adopted a pre-existing tool that had not been previously assessed for validity and reliability. We based the initial item generation process on literature review findings, and we conducted the pilot test with a small sample—without assessment for construct validity. Recognition of these limitations was the primary motivation to validate the tool further using quantitative methods.

The second limitation is that we did not apply other forms of reliability and validity, such as alternate forms reliability, test–retest reliability, and criterion-related validity. We distributed a single version of the survey to hospital staff only once before we conducted the education program. Without distributing an alternate survey that worded questions differently, we could not assess alternate forms reliability. With a single round of baseline data, we could not examine the tool’s temporal stability. We conducted a literature search but could not find an existing “gold standard” scale for measuring TIC KAP for healthcare workers or workers in other professional sectors. Without a “gold standard” scale to compare against, we could not assess criterion-related validity.³¹

A third limitation is the generalizability of the survey. The primary objective of this study is to test the theorized item-factor relationships using CFA, for which the sample size is large enough. However, the sample is drawn exclusively from a single institution, and the response rate (22.3%) is low compared with typical online surveys.³² Although there was diversity in the role and department affiliation, the possibility of bias exists. To make the scale truly generalizable for the field, we recommend that additional data be collected through a multicenter study approach from other healthcare agencies and institutions for further validation. Also, this study was done only in a pediatric institution. The original literature on ACE exposure was done in an adult cohort,³ and therefore, it is well known that this science is relevant to adult populations as well. Further testing of this tool for applicability in adult healthcare institutions is needed because this work is of equal importance in the adult healthcare field.

Finally, goodness of fit indices CFI and TLI came close but still fell short of the commonly accepted threshold for good model fit. However, experts have recommended a balanced approach to consider both the theory and model fit. Too much focus on adhering to cutoffs of goodness of fit indices could lead to type 1 error, incorrectly rejecting an acceptable model.^{23,33} In other words, theory and the RMSEA suggest that this is an acceptable model. It would be incorrect to reject it solely due to CFI and TLI values missing the commonly accepted thresholds. Having considered both the theory and the model fit, we believe our scale will be useful for implementing TIC in healthcare

settings despite not meeting all of the goodness of fit indices.

CONCLUSIONS

We believe that healthcare organizations can take a proactive approach to improve the health and well-being of their patients by implementing a trauma-informed approach to minimize the impact of childhood trauma and adversity. Accurately assessing the needs of learners is an essential step in transforming the KAP among healthcare organizations and healthcare professionals as they relate to TIC. This tool will allow pediatric organizations to identify gaps in KAP among staff to subsequently develop strategies to achieve a culture of TIP.

DISCLOSURE

The authors have no financial interest to declare in relation to the content of this article.

REFERENCES

- Marsac ML, Kassam-Adams N, Hildenbrand AK, et al. Implementing a trauma-informed approach in pediatric health care networks. *JAMA Pediatr.* 2016;170:70–77.
- Substance Abuse and Mental Health Services Administration. *SAMHSA’s Concept of Trauma and Guidance for a Trauma-Informed Approach* (HHS Publication No. SMA 14-4884). Rockville, MD: Substance Abuse and Mental Health Services Administration; 2014.
- Boles J. Trauma-informed care: an intentional approach. *Pediatric Nursing.* 2017;43(5):250.
- Felitti VJ, Anda RF, Nordenberg D, et al. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) Study. *Am J Prev Med.* 1998;14:245–258.
- Shonkoff JP, Garner AS, Siegel BS, et al. Committee on psychosocial aspects of child and family health; committee on early childhood, adoption, and dependent care; section on developmental and behavioral pediatrics. The lifelong effects of early childhood adversity and toxic stress. *Pediatrics.* 2012;129:e232–e246. doi:10.1542/peds.2011-2663
- Leitch L. Action steps using ACEs and trauma-informed care: a resilience model. *Health Justice.* 2017;5:5.
- Burke Harris N, Silvério Marques S, Oh D, et al. Prevent, screen, heal: collective action to fight the toxic effects of early life adversity. *Acad Pediatr.* 2017;17(7S):S14–S15.
- Koetting C. Trauma-informed care: helping patients with a painful past. *J Christ Nurs.* 2016;33:206–213.
- Morrissey JP, Jackson EW, Ellis AR, et al. Twelve-month outcomes of trauma-informed interventions for women with co-occurring disorders. *Psychiatr Serv.* 2005;56:1213–1222.
- Suarez E, Jackson DS, Slavin LA, et al. Project Kealahou: improving Hawaii’s system of care for at-risk girls and young women through gender-responsive, trauma-informed care. *Hawaii J Med Public Health.* 2014;73:387–392.
- American Academy of Pediatrics. Trauma Toolbox for Primary Care. 2014. <https://www.aap.org/en-us/advocacy-and-policy/aap-health-initiatives/healthy-foster-care-america/Pages/Trauma-Guide.aspx>. Accessed October 31, 2018.
- Menschner C, Maul A. Key Ingredients for Successful Trauma-Informed Care Implementation. Center for Health Care Strategies, INC; 2016. <https://www.chcs.org/media/Brief-Key-Ingredients-for-TIC-Implementation-1.pdf>. Accessed October 31, 2018.
- Kerker BD, Storfer-Isser A, Szilagyi M, et al. Do pediatricians ask about adverse childhood experiences in pediatric primary care? *Acad Pediatr.* 2016;16:154–160.

14. Data Resource Center for Child & Adolescent Health. The National Survey of Children's Health. <http://childhealthdata.org/learn/NSCH>. Accessed October 31, 2018.
15. Pilcher J. Learning needs assessment: not only for continuing education. *J Nurses Prof Dev*. 2016;32:122–129. doi:10.1097/NND.0000000000000245
16. Schrader PG, Lawless KA. The Knowledge, Attitudes, and Behaviors Approach: How to evaluate performance and learning in complex environments. *Perform Improv*. 2004;43:8–15.
17. Stokes Y, Jacob JD, Gifford W, et al. Exploring nurses' knowledge and experiences related to trauma-informed care. *Glob Qual Nurs Res*. 2017;4:1–10.
18. Bruce MM, Kassam-Adams N, Rogers M, et al. Trauma providers' knowledge, views, and practice of trauma-informed care. *J Trauma Nurs*. 2018;25:131–138.
19. Baker C, Brown S, Wilcox P, Overstreet S, Arora P. Development and psychometric evaluation of the Attitudes Related to Trauma-Informed Care (ARTIC) Scale. *School Ment Health*. 2016;8:61–76. doi:10.1007/s12310-015-9161-0
20. Abdoh N, Bernardi E, McCarthy A. Knowledge, Attitudes, and Practices of Trauma-Informed Practice: A Survey of Health Care Professionals and Support Staff at Alexander Street Community. Unpublished paper, 2017. <https://open.library.ubc.ca/media/download/pdf/52966/1.0343062/5>. Accessed October 31, 2018.
21. Jackson DL, Gillaspay JA, Purc-Stephenson R. Reporting practices in confirmatory factor analysis: an overview and some recommendations. *Psychol Methods*. 2009;14:6–23.
22. Kenny DA. Measuring Model Fit. Updated 2015. <http://davidakenny.net/cm/fit.htm>. Accessed May 24, 2019.
23. Hooper D, Coughlan J, Mullen MR. Structural equation modelling: guidelines for determining model fit. *Electron J Bus Res Methods*. 2008;6:53–60.
24. Bentler PM. Comparative fit indexes in structural models. *Psychol Bull*. 1990;107:238–246.
25. Bentler PM, Bonnet DC. Significance tests and goodness of fit in the analysis of covariance structures. *Psychol Bull*. 1980;88:588–606.
26. Kenny DA. Respecification of Latent Variable Models. 2011. <http://davidakenny.net/cm/respec.htm>. Accessed July 29, 2019.
27. Brown TA. *Confirmatory Factor Analysis for Applied Research*. New York, NY: The Guilford Press; 2006.
28. MacCallum R, Browne M, Sugawara H. Power analysis and determination of sample size for covariance structure modeling. *Psychol Methods*. 1996;1:130–149. doi:10.1037/1082-989X.1.2.130
29. Myers ND, Ahn S, Jin Y. Sample size and power estimates for a confirmatory factor analytic model in exercise and sport: a Monte Carlo approach. *Res Q Exerc Sport*. 2011;82:412–423.
30. Tavakol M, Dennick R. Making sense of Cronbach's alpha. *Int J Med Educ*. 2011;2:53–55.
31. DeVellis R. *Scale Development*. Los Angeles, CA: Sage; 2017.
32. Nulty D. The adequacy of response rates to online and paper surveys: what can be done? *Assess Eval High Edu*. 2008;33:301–314.
33. Marsh HW, Hau KT, Wen Z. In search of golden rules: comment on hypothesis-testing approaches to setting cutoff values for fit indexes and dangers in overgeneralizing Hu and Bentler's findings. *Struct Equ Modeling*. 2004;11:320–341.